

Liberty Hospital participates in the Cancer Program Practice Profile Reports (CP3R) as part of their participation in Commission on Cancer accreditation. Data on breast cancer from 2013 through 2015 are reported here. Liberty Hospital met the benchmark for the quality care measures below. In 2015, in three of the four criteria that are benchmarked, Liberty Hospital ranked higher than the Missouri and national accredited hospitals. The Cancer Committee at Liberty Hospital continually monitors our data to ensure we are providing quality care.

<b>BREAST</b>	<b>2013 Liberty Hospital Performance Rate (EPR)</b>	<b>2014 Liberty Hospital Performance Rate (EPR)</b>	<b>2015 Liberty Hospital Performance Rate (EPR)</b>	<b>COC Standard &amp; Benchmark</b>	<b>2015 Comparison - Missouri CoC accredited hospitals</b>	<b>2015 Comparison- U.S. CoC accredited hospitals (CCCP category)</b>
<b>1. Radiation is administered within 1 year (365 days) of diagnosis for women under age 70 receiving breast conservation surgery for breast cancer (BCSRT)</b>	100%	100%	100%	Standard 4.4 90%	95.1%	90.2%
<b>2. Tamoxifen or third generation aromatase inhibitor is recommended or administered within 1 year (365 days) of diagnosis for women with AJCC T1c or Stage IB - III hormone receptor positive breast cancer (HT)</b>	86.7%	97.1%	100%	Standard 4.4 90%	94%	90.1%
<b>3. Radiation therapy is recommended or administered following any mastectomy within 1 year (365 days) of diagnosis of breast cancer for women with &gt;= 4 positive regional lymph nodes (MASTRT)</b>	66.7%	100%	100%	Standard 4.4 90%	91.4%	84.1%
<b>4. Combination chemotherapy is recommended or administered within 4 months (120 days) of diagnosis for women under 70 with AJCC T1cN0, or Stage IB - III hormone receptor negative breast cancer (MAC)</b>	100%	100%	100%	N/A	94.6%	92.5%